



A database for post-genome analysis

When the Human Genome Project was initiated in the late 1980s, it was perceived as the ultimate project to uncover the blueprint of life. Although the goal of sequencing the entire 3.0 billion base pairs of the human genome by 2005 is likely to be achieved, whether we will have the blueprint of life at that time is quite questionable.

GENETWORK

First of all, as we have learned from the complete genomes of yeast and several bacteria, the biological function of a large fraction of the genes is found to be either unknown or uncharacterized. Secondly, and more importantly, because the genes and gene products are only the individual components that make up a biological system, the understanding of how each component works is not sufficient to understand the entire system. The post-genomic analysis, as we define it here, includes both experimental and informatics approaches to uncover systematically the interactions and pathways of genes and molecules, which can be considered as the wiring diagrams of the biological system. The complete catalog of components and the complete catalog of wiring diagrams together can be called the blueprint of life.

KEGG (Kyoto Encyclopedia of Genes and Genomes) is an informatics project for the post-genomic analysis, which was initiated in 1995 under the Human Genome Program of the Ministry of Education, Science, Sports and Culture in Japan. Its objectives are threefold: (1) to compile the current knowledge of molecular pathways

and genetic pathways; (2) to experimentally observation of genetic, biochemical, and molecular and cellular biology. In the past two years, KEGG contained only the metabolic pathways, but starting in July 1997 a number of regulatory pathways, such as signal transduction, cell cycle and developmental pathways, are being placed online. (2) KEGG maintains the gene catalog of every organism that has been sequenced, and each component in the catalog is to be mapped on to the KEGG pathways. (3) In addition to these database efforts, KEGG aims at developing new informatics technologies that are associated with interactions and pathways.

KEGG is a part of the Japanese GenomeNet WWW server¹ and is linked to all the major molecular biology databases by the ENRICH (Enrichment) system. Figure 1 shows a portion of the KEGG metabolic pathway diagram for phenylalanine, tyrosine, and tryptophan biosynthesis, where each box represents an enzyme with the EC number inside. The box is clickable to retrieve the corresponding enzyme entry of the EMBL database², which is the starting point of retrieving related entries of chemical compounds, molecular structures,




Figure 1. Examples of using KEGG at <http://www.genome.ad.jp/kegg>. (a) From the KEGG table of contents, choose the molecular catalog. Enzymes by SCOP 3D-fold; select alpha and beta in B1, copy all the description in the category of alpha beta (C14M1) and, select the 'Pathway' option, paste in the search box, and search against '3D structures in PDB'. Several pathways, each of which contains at least one enzyme with a TIM barrel according to SCOP, will be indicated in the result screen. The view shown is (run of p080100) phenylalanine, tyrosine and tryptophan biosynthesis. The blue boxes are the enzymes whose three-dimensional structures are known and those marked with red are the ones found to correspond to the SCOP classification. (b) From the KEGG table of contents, choose the catalog of *Escherichia coli* operons, copy all the description in the *trp* operon, select the 'Pathway' option, paste in the search box, and search against *Escherichia coli*. Select the 'second set' (phenylalanine, tyrosine and tryptophan biosynthesis) pathway. The green boxes are the enzymes whose genes exist and those marked with red are the ones found to correspond to the *trp* operon.

GENETWORK

three-dimensional structures, and genetic diseases among others. KEGG contains structural and functional classifications of molecules and genes in the form of what we call hierarchical trees, in which the headings and subheadings are clickable to unfold or fold branches. Figure 1(a) is a result (treated in red) of matching the Jia (TAM) based pictures in the hierarchical table derived from the SCOP database with the KEGG metabolic pathway diagrams where the enzymes with known PDB (Protein Data Bank) structures are shown in blue boxes. This indicates possible gene duplications in the formation of the tryptophan biosynthetic pathway⁶.

One of the most unique aspects of KEGG is the automatic generation of organism-specific pathways by matching the gene catalogs being produced by the genome sequencing projects and the reference pathway diagrams manually drawn and updated. In Figure 1(b) the enzymes colored in green indicate that the corresponding genes are found in the Echerichia coli gene catalog. Those marked in red belong to the tryptophan operon, and the genome map section of KEGG can also be utilized with a java-compatible browser, for example, to see any correlation between the physical proximity of genes

in the genome and the functional proximity of gene products in the pathway. An important consequence of mapping gene products on the pathway diagrams is the validation of the initial gene assignments. In case the pathway is not continuous because of missing gene products, KEGG provides computational tools to assist re-examination of gene function assignment⁷ and further analysis of possible existence of alternative paths⁸.

While KEGG tries to cover a diverse range of pathways at a high level of abstraction, there are complementary resources that contain more detailed data and knowledge in specific pathways. We have started collaboration with WIT (ref. 9) for metabolic pathways and are open to any other collaborations. The mirror sites of KEGG are being established in the USA and UK (ref. 10). In addition to the Internet version, KEGG is available in CD-ROM for Macintosh and Windows where pathway diagrams, hierarchical texts, and genome maps are all to be handled by a java-compatible browser. The content of CD-ROM can be downloaded by anonymous FTP (ref. 11) and used in UNIX as well. We also plan to start distributing the KEGG server to be mirrored in a local environment.

Minoru Kanehisa

kanehisa@kaiyaku.kyoto-u.ac.jp
<http://www.kaiyaku.kyoto-u.ac.jp/kanehisa/>

Institute for Chemical Research, Kyoto University, Uji, Kyoto 611, Japan

References

- 1 Goto, S. et al. (1996) Pacific Symposium on Biocomputing 1997, pp. 175-186
- 2 <http://www.genome.ad.jp>
- 3 <http://www.genome.ad.jp/digest/digestlinks.html>
- 4 http://www.genome.ad.jp/htbin/show_man/figand
- 5 Morin, A.G., Brenner, S.E., Hubbard, T. and Chothia, C. (1995) *J. Mol. Biol.* 247, 535-540
- 6 Widmann, M. and Eisenberg, D. (1993) *Proc. Natl. Acad. Sci. U. S. A.* 90, 1379-1383
- 7 <http://www.genome.ad.jp/kegg/comp/GIT.html>
- 8 <http://www.genome.ad.jp/kegg/comp/pathcomp.html>
- 9 <http://www.cmc.mcgill.ca/wit/>
- 10 <http://kegg.com>
- 11 <http://kegg.genome.ad.jp/CD/>

TECHNICAL TIPS ONLINE

<http://www.elsevier.com/locate/tio> <http://www.elsevier.nl/locate/tio> (In Europe)

Technical Tips Online publishes short, peer-reviewed, molecular biology techniques articles in a Web-based environment. The articles describe novel methods or significant improvements to existing methods in any aspect of molecular biology.

New Technical Tip articles published recently in **Technical Tips Online** include:

Choi, C., Mikes, D.G. and Pritikin, F. (1997) A PCR method for distinguishing cells from mouse strains 129 and C57BL/6 for gene knockout studies. *Technical Tips Online* (<http://www.elsevier.com/locate/tio>) T01139

Granger, B.L. (1997) A glass bead method for picking bacterial colonies. *Technical Tips Online* (<http://www.elsevier.com/locate/tio>) T01175

Heflich, S., Nairn, F., Lemmon, C. and Bonacc, J. (1997) There is still life after death for T-cell line studies. <http://www.elsevier.com/locate/tio> T01186

Lin, D. and Gachem, N. (1997) Use of a simplified and rapid site screen protocol for the detection of recombinant plasmids. *Technical Tips Online* (<http://www.elsevier.com/locate/tio>) T01172

Loth, A., Im, Y.H., Sengupta, P. and Hanson, R. (1997) A method for preparing DM labeled tissue that allows single-cell resolution with conventional epifluorescence microscopy. <http://www.elsevier.com/locate/tio> T01119

Menck, J. (1997) Two years Echerichia coli/plasmid vector designed for yeast one- and two-hybrid screens that allow directional cDNA cloning. <http://www.elsevier.com/locate/tio> T01111

Parkinson, N., Barclay, J., Gordon, M. and Res, H. (1997) An inverse PCR strategy for the recovery of end fragments from the pRML YAC vector. *Technical Tips Online* (<http://www.elsevier.com/locate/tio>) T01160

Schmitt, H. (1997) Rapid high-throughput partitioning of genomic DNA from mouse and rat tails for use in transgenic testing. *Technical Tips Online* (<http://www.elsevier.com/locate/tio>) T01146

Spielman, P., Van Gels, M., Mervens, L. and Van der, N. (1997) Improved protocol for chromatin fibers from blood cells. <http://www.elsevier.com/locate/tio> T01123

Spradling, B., Berg, H.H. and Myers, P.B. (1997) Recycling cells in PCR and immunology studies. <http://www.elsevier.com/locate/tio> T01090

Editor: Adrian Brier, Institute for Cell and Molecular Biology at the University of Edinburgh